

Radiative Processes

Simon Casassus


Astronomía, Universidad de Chile

<http://www.das.uchile.cl/~simon>

- I Radiative Transfer
- II Electromagnetic wave propagation
- III Radiation
- IV Scattering and Diffraction
- V Free-free, Synchrotron and Compton Scattering
- VI Radiative Transitions

Part V

Free-free, Synchrotron and Compton Scattering



Relativity


Quadrivectors
Covariance in
electrodynamics

Outline

1 Relativity

Quadrivectors

Covariance in electrodynamics



Relativity

Quadrivectors


Covariance in
electrodynamics

Outline

1 Relativity

Quadrivectors

Covariance in electrodynamics



Relativity

Quadrivectors

Covariance in
electrodynamics


1.1- Quadrivectors

In this section we switch to CGS units, better adapted to describe the symmetry between \vec{E} and \vec{B} (bibliography: Rybicki & Lightman).

- We define $x^\mu = (ct, x, y, z)$ as the contravariant position quadrivector, whose norm is $s^2 = \eta_{\mu\nu} x^\mu x^\nu$ (using the implicit sum notation).
- We also introduce $x_\mu = (-ct, x, y, z)$ as the covariant position quadrivector.
- $x_\mu = \eta_{\mu\nu} x^\nu$, with

$$\eta_{\mu\nu} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \quad (1)$$

- With $\eta^{\mu\nu} \equiv \eta_{\mu\nu}$, we have $x^\mu = \eta^{\mu\nu} x_\nu$.
- Note that $\eta^{\mu\sigma} \eta_{\sigma\nu} = \delta_\nu^\mu$.



1.1- Quadrivectors

- We change reference system from \mathcal{S} to \mathcal{S}' , in uniform translation with velocity v towards \hat{x} relative to \mathcal{S} .
- A contravariant 4V transforms as

$$x'^{\mu} = \Lambda^{\mu}_{\nu} x^{\nu}, \text{ where } \Lambda^{\mu}_{\nu} = \begin{bmatrix} \gamma & -\beta\gamma & 0 & 0 \\ -\beta\gamma & \gamma & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad (2)$$


with $\beta = v/c$, and $\gamma = 1/\sqrt{1 - v^2/c^2}$.

- For a covariant 4V (tarea),

$$x'_{\mu} = \tilde{\Lambda}_{\mu}^{\nu} x_{\nu}, \text{ with } \tilde{\Lambda}_{\mu}^{\nu} = \eta_{\mu\tau} \Lambda^{\tau}_{\sigma} \eta^{\sigma\nu}. \quad (3)$$

- $\tilde{\Lambda}_{\mu}^{\nu}$ is the inverse of Λ^{μ}_{ν} :

$$\Lambda^{\sigma}_{\nu} \tilde{\Lambda}_{\sigma}^{\mu} = \delta_{\nu}^{\mu}, \text{ and } \tilde{\Lambda}_{\mu}^{\alpha} x'^{\mu} = x^{\alpha}. \quad (4)$$



Relativity

Quadrivectors

Covariance in
electrodynamics

1.1- Quadriectors

- The product of two 4Vs A^μ and B_μ is Lorentz invariant:


$$A^\mu B_\mu = A'^\mu B'_\mu. \quad (5)$$

- We also have the velocity 4V, $U^\mu \equiv \frac{dx^\mu}{d\tau}$, in which $d\tau$ is the relativistic interval. (i.e. of proper time) between x^μ and $x^\mu + dx^\mu$.
- In components (tarea), $U^\mu = \gamma_u(c, \vec{u})$, with $\gamma_u = 1/\sqrt{1 - u^2/c^2}$.
- If we change to $U' = \Lambda^\mu{}_\nu U^\nu$,


$$\gamma_{u'} = \gamma \gamma_u (1 - \frac{uv}{c^2} \cos(\theta)), \quad \text{with } \theta = \angle(\vec{u}, \vec{v}). \quad (6)$$

- In the system that is bound to a particle with velocity \vec{u} , $U' = c(1, \vec{0})$, for a 4V A^μ ,

$$A'^0 = -\frac{1}{c} U^\mu A_\mu = -\frac{1}{c} U'^\mu A'_\mu \dots \quad (7)$$



1.1- Quadrivectors



Relativity

Quadrivectors

Covariance in
electrodynamics

- We note that the phase of a plane wave must be Lorentz invariant because the simultaneous cancellation of \vec{E} and \vec{B} in one system implies their cancellation in any other system.
- Let's introduce $k^\mu = (\omega/c, \vec{k})$:

$$k^\mu x_\mu = \vec{k} \cdot \vec{x} - \omega t = \text{invariant} \Rightarrow k^\mu \text{ is 4V.} \quad (8)$$

- We can use Eq. 7 to deduce the relativistic Doppler effect (tarea)

$$ck'^0 = \omega' = -U^\mu k_\mu = \omega \gamma \left(1 - \frac{v}{c} \cos(\theta)\right). \quad (9)$$

1.1- Quadrivectors

- The gradient operator is another example of 4V. If λ is a scalar invariant, then

$$\lambda_{,\mu} \equiv \frac{\partial \lambda}{\partial x^\mu} \text{ is a covariant 4V, and} \quad (10)$$

$$\lambda^{,\mu} \equiv \frac{\partial \lambda}{\partial x_\mu} \text{ is a contravariant 4V.} \quad (11)$$


- Proof: from $x^\nu = \tilde{\Lambda}^\nu{}_\mu x'^\mu$, we have that $\frac{\partial x^\nu}{\partial x'^\mu} = \tilde{\Lambda}^\nu{}_\mu$, and since $\lambda' = \lambda$,

$$\lambda'_{,\mu} = \frac{\partial x^\nu}{\partial x'^\mu} \frac{\partial \lambda}{\partial x^\nu}. \quad (12)$$

- We extend the properties of 4Vs to tensors in general: a tensor of order n transforms as the product of n 4Vs.
- For example,

$$T'^{\mu\nu} = \Lambda^\mu{}_\sigma \Lambda^\nu{}_\tau T^{\sigma\tau},$$

$$T'^\mu{}_\nu = \Lambda^\mu{}_\sigma \tilde{\Lambda}^\tau{}_\nu T^\sigma{}_\tau.$$




Outline

1 Relativity

Quadrivectors

Covariance in electrodynamics



Relativity

Quadrivectors

Covariance in
electrodynamics

1.2- Covariance in electrodynamics

- Charge conservation, $\frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot \vec{J} = 0$, can be written as

$$J^{\mu}_{,\mu} = 0, \text{ using the four-current } J^{\mu} = (\rho c, \vec{J}). \quad (13)$$

- In the Lorentz Gauge, and using CGS units,


$$\nabla^2 \vec{A} - \frac{1}{c^2} \frac{\partial^2 \vec{A}}{\partial t^2} = -\frac{4\pi}{c} \vec{J} = \partial_{\alpha} \partial^{\alpha} \vec{A}, \quad (14)$$

$$\nabla^2 \Phi - \frac{1}{c^2} \frac{\partial^2 \Phi}{\partial t^2} = -4\pi \rho = \partial_{\alpha} \partial^{\alpha} \Phi. \quad (15)$$

- With $A^{\mu} = (\Phi, \vec{A})$,

$$A^{\beta,\alpha}_{,\alpha} = -\frac{4\pi}{c} J^{\beta}, \text{ in which } A^{\beta,\alpha}_{,\alpha} = \frac{\partial^2}{\partial x_{\alpha} \partial x^{\alpha}} A^{\beta}. \quad (16)$$

- The Lorentz gauge $\vec{\nabla} \cdot \vec{A} + \frac{1}{c} \frac{\partial \Phi}{\partial t} = 0$ can be written simply as $A^{\alpha}_{,\alpha} = 0$.



Relativity

Quadrivectors

Covariance in
electrodynamics

1.2- Covariance in electrodynamics

- In order to write the Maxwell equations in their covariant form, we introduce the field tensor

$$F_{\mu\nu} = A_{\nu,\mu} - A_{\mu,\nu}. \quad (17)$$

- With $\vec{B} = \vec{\nabla} \times \vec{A}$ and $\vec{E} = -\vec{\nabla}\Phi - \frac{1}{c} \frac{\partial \vec{A}}{\partial t}$ (tarea): 2


$$F_{\mu\nu} = \begin{bmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & B_z & -B_y \\ E_y & -B_z & 0 & B_x \\ E_z & B_y & -B_x & 0 \end{bmatrix} \quad (18)$$

- The Maxwell equations $\vec{\nabla} \cdot \vec{E} = 4\pi\rho$ and $\vec{\nabla} \times \vec{B} - \frac{1}{c} \frac{\partial \vec{E}}{\partial t} = \frac{4\pi}{c} \vec{J}$ can be written (tarea)

$$F_{\mu\nu}{}^{,\nu} = \frac{4\pi}{c} J_{\mu}. \quad (19)$$

- The 'internal' equations $\vec{\nabla} \cdot \vec{B} = 0$ and $\vec{\nabla} \times \vec{E} + \frac{1}{c} \frac{\partial \vec{B}}{\partial t} = 0$ are written (tarea)

$$F_{\mu\nu,\sigma} + F_{\sigma\mu,\nu} + F_{\nu\sigma,\mu} = 0. \quad (20)$$




Relativity

Quadrivectors

Covariance in
electrodynamics

1.2- Covariance in electrodynamics



Relativity

Quadrivectors

Covariance in
electrodynamics

- We use the covariance of $F_{\mu\nu}$ to infer the transformation laws for the fields \vec{E} y \vec{B} :

$$F'_{\mu\nu} = \tilde{\Lambda}_{\mu}^{\alpha} \tilde{\Lambda}_{\nu}^{\beta} F_{\alpha\beta}. \quad (21)$$

- In terms of components we get (tarea):

$$\begin{aligned} E'_{\parallel} &= E_{\parallel}, & B'_{\parallel} &= B_{\parallel}, \\ E'_{\perp} &= \gamma(\vec{E}_{\perp} + \vec{\beta} \times \vec{B}), & B'_{\perp} &= \gamma(\vec{B}_{\perp} + \vec{\beta} \times \vec{E}). \end{aligned} \quad (22)$$

- We see that \vec{E} and \vec{B} get mixed up, and if $\vec{B} = 0$ in S , then when changing to S' we have $\vec{B}' \neq 0$.

1.2- Covariance in electrodynamics

- In order to extend the Lorentz force, we introduce the momentum quadrivector (four-momentum) $P^\mu = m_o U^\beta$, where m_o is the rest mass. We write $P^\mu = (E/c, \vec{P})$, in which E is the total energy of the particle (which is $E = m_o c^2$ at rest).
- The acceleration 4V (four-acceleration) is

$$a^\mu = \frac{dU^\mu}{d\tau}, \quad (23)$$


and in order to recover Newton's 2nd law in the non-relativistic limit, the four-force must be

$$F^\mu = m_o a^\mu = \frac{dP^\mu}{d\tau}. \quad (24)$$

- We write the 4-Lorentz force with

$$F^\mu = \frac{q}{c} F^\mu{}_\nu U^\nu. \quad (25)$$

- in components, (tarea) $\vec{F} = q(\frac{\vec{v}}{c} \times \vec{B}) + q\vec{E}$.



Relativity

Quadrivectors

Covariance in
electrodynamics