

# Radiative Processes

Simon Casassus


Astronomía, Universidad de Chile

<http://www.das.uchile.cl/~simon>

- I Radiative Transfer
- II Electromagnetic wave propagation
- III Radiation
- IV Scattering and Diffraction
- V Free-free, Synchrotron and Compton Scattering
- VI Radiative Transitions

# Part III

## Radiation



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials

Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

### Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

### Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

### Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

### Non-relativistic applications

# 1- Green function for the wave equation

- In order to determine  $\vec{A}(\vec{x}, t)$  and  $\Phi(\vec{x}, t)$ , we need to solve the wave equation with source terms. For a generic field  $\Psi(\vec{x}, t)$ ,

$$\nabla^2 \Psi - \frac{1}{c^2} \frac{\partial^2 \Psi}{\partial t^2} = -4\pi f(\vec{x}, t). \quad (1)$$

- It is convenient to use the Fourier time-domain,


$$\psi(\vec{x}, t) = \frac{1}{2\pi} \int_{-\infty}^{+\infty} \psi(\vec{x}, \omega) e^{-i\omega t} d\omega, \quad (2)$$

whose inverse is

$$\psi(\vec{x}, \omega) = \int_{-\infty}^{+\infty} \psi(\vec{x}, t) e^{i\omega t} dt. \quad (3)$$

- Injecting Ec. 2 in Ec. 1, we reach the Helmholtz equation:

$$(\nabla^2 + k^2) \Psi(\vec{x}, \omega) = -4\pi f(\vec{x}, \omega). \quad (4)$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

## 1- Green function for the wave equation

- The Helmholtz equation Eq. 4 is very similar to the Poisson equation, and we can anticipate the use of similar machinery in its solution. The Green function satisfies

$$(\nabla^2 + k^2)G_k(\vec{x}, \vec{x}') = -4\pi\delta(\vec{x}, \vec{x}'). \quad (5)$$


- Changing coordinates to a system with origin in  $\vec{x}'$ , we see that Eq. 5 has spherical symmetry, and  $G_k(\vec{x}, \vec{x}') = G_k(R)$ , with  $R = |\vec{R}|$  and  $\vec{R} = \vec{x} - \vec{x}'$
- Eq. 5 can thus be written as

$$\frac{1}{R} \frac{d^2}{dR^2} (R G_k) + k^2 G_k = -4\pi\delta(\vec{R}). \quad (6)$$

- If  $R \neq 0$ , the solution to Eq. 6 is  $R G_k = A e^{ikR} + B e^{-ikR}$ , where the constants  $A$  and  $B$  do not depend on  $k$ . In order to determine these constants, we use the case  $k = 0$ , i.e. Poisson, whose solution is  $G_{k=0}(R) = 1/R$ ,  $\rightarrow A + B = 1$ .
- Thus the general solution to Eq. 6 is

$$G_k(R) = A G_k^+(R) + B G_k^-(R), \text{ with } G_k^\pm = \frac{e^{\pm ikR}}{R}$$

$$\text{and } A + B = 1. \quad (7)$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

# 1- Green function for the wave equation

- The  $A$  and  $B$  values depend on the initial conditions, i.e. on the boundary conditions in time. To see this, we return to the time domain and we generalize Eq. 5:

$$\left(\nabla^2 - \frac{1}{c^2} \frac{\partial}{\partial t^2}\right) G^\pm(\vec{x}, t; \vec{x}', t') = -4\pi\delta(\vec{x} - \vec{x}')\delta(t - t'). \quad (8)$$

- Now, returning to the frequency domain  $\omega$ , we generalize Eq. 5 to


$$(\nabla^2 + k^2)G_k(\vec{x}, \vec{x}'; t') = -4\pi\delta(\vec{x}, \vec{x}')e^{i\omega t'}, \quad (9)$$

with solution  $G_k^\pm(R)e^{i\omega t'}$ .

- To return once more to the time domain, we use Eq. 2, and

$$G^\pm(R; t, t') = G^\pm(R, \tau) = \frac{1}{2\pi} \int \frac{e^{\pm ikR - i\omega\tau}}{R} d\omega,$$

where  $\tau = t - t'$ .



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

## 1- Green function for the wave equation

- For a non-dispersive medium (one with  $\omega/k = c$ ), we reach

$$G^{\pm}(\vec{x}, t; \vec{x}', t') = \frac{\delta\left(t' - \left[t \mp \frac{|\vec{x} - \vec{x}'|}{c}\right]\right)}{|\vec{x} - \vec{x}'|} \quad (10)$$

- We apply this Green function to write the generic solutions of Ec. 1:


$$\Psi^{\pm}(\vec{x}, t) = \int d^3x' dt' G^{\pm}(\vec{x}, t; \vec{x}', t') f(\vec{x}', t'). \quad (11)$$

- The  $+$  case corresponds to the retarded solution, with an entry or an initial condition  $\psi_{\text{in}}$  (valid before the sources  $f$  are activated, at  $t = 0$ , so  $f(\vec{x}, t) = 0$  if  $t < 0$ ):

$$\Psi^{+}(\vec{x}, t) = \Psi_{\text{in}}(\vec{x}, t) + \int d^3x' dt' G^{+}(\vec{x}, t; \vec{x}', t') f(\vec{x}', t'), \quad (12)$$

where we see that if  $t < 0$ , there is no  $\vec{x}$  for any given  $\vec{x}'$  such that

$\left[t - \frac{|\vec{x} - \vec{x}'|}{c}\right] > 0$ . Hence if  $t < 0$ ,  $\int d^3x' dt' G^{+}(\vec{x}, t; \vec{x}', t') f(\vec{x}', t') = 0$ , and  $\Psi(\vec{x}, t) = \Psi_{\text{in}}(\vec{x}, t)$ .



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications


# 1- Green function for the wave equation

- Instead the  $-$  case corresponds to the anticipated solution, with an exit condition  $\Psi_{\text{out}}$  (after the sources  $f$  are deactivated, at  $t = 0$ , so  $f(\vec{x}, t) = 0$  if  $t > 0$ ),

$$\Psi^{-}(\vec{x}, t) = \Psi_{\text{out}}(\vec{x}, t) + \int d^3x' dt' G^{-}(\vec{x}, t; \vec{x}', t') f(\vec{x}', t'), \quad (13)$$

where we see that if  $t > 0$ , there is no  $\vec{x}$  for any given  $\vec{x}'$  such that  $\left[ t + \frac{|\vec{x} - \vec{x}'|}{c} \right] < 0$ . Hence if  $t > 0$ ,  $\int d^3x' dt' G^{-}(\vec{x}, t; \vec{x}', t') f(\vec{x}', t') = 0$ , and  $\Psi(\vec{x}, t) = \Psi_{\text{out}}(\vec{x}, t)$ .

- In general we use the retarded solution  $\Psi^{+}$ .



## Green function for the wave equation

### Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

### Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

### Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

### Non-relativistic applications



# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

### Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

### Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

### Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

### Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials

Retarded electromagnetic field

## 3 Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation


## 4 Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials

Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications

## 2.1- Application of the Green function to the electrodynamic potentials

- We normally use the retarded solution, with initial condition  $\Psi_{\text{in}} = 0$ , or, in compact notation,


$$\Psi(\vec{x}, t) = \int d^3x' \frac{[f(\vec{x}', t')]_{\text{ret}}}{|\vec{x} - \vec{x}'|}, \quad (14)$$

where  $[(\dots)]_{\text{re}}$  means to evaluate in  $t' = t - |\vec{x} - \vec{x}'|/c$ .

- Applying to the electrodynamic potentials,

$$\Phi(\vec{x}, t) = \frac{1}{4\pi\epsilon_0} \int d^3x' \frac{[\rho(\vec{x}', t')]_{\text{ret}}}{|\vec{x} - \vec{x}'|}, \quad (15)$$

$$\vec{A}(\vec{x}, t) = \frac{\mu_0}{4\pi} \int d^3x' \frac{[\vec{J}(\vec{x}', t')]_{\text{ret}}}{|\vec{x} - \vec{x}'|}. \quad (16)$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials

Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials

Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications

## 2.2- Retarded electromagnetic field

- In order to calculate  $\vec{E}$  and  $\vec{B}$ , we use  $\vec{B} = \vec{\nabla} \times \vec{A}$  and  $\vec{E} = -\vec{\nabla}\phi - \frac{\partial \vec{A}}{\partial t}$ .
- Alternatively, we can use the Maxwell equations to reach:


$$\nabla^2 \vec{E} - \frac{1}{c^2} \frac{\partial^2 \vec{E}}{\partial t^2} = -\frac{1}{\epsilon_0} \left( -\vec{\nabla} \rho - \frac{1}{c^2} \frac{\partial \vec{J}}{\partial t} \right), \quad (17)$$

$$\nabla^2 \vec{B} - \frac{1}{c^2} \frac{\partial^2 \vec{B}}{\partial t^2} = -\mu_0 \vec{\nabla} \times \vec{J}. \quad (18)$$

- Using the Green function, we get

$$\vec{E}(\vec{x}, t) = \frac{1}{4\pi\epsilon_0} \int d^3x' \frac{1}{R} \left[ -\vec{\nabla}' \rho - \frac{1}{c^2} \frac{\partial \vec{J}}{\partial t'} \right]_{\text{ret}}, \quad (19)$$

$$\vec{B}(\vec{x}, t) = \frac{\mu_0}{4\pi} \int d^3x' \frac{1}{R} \left[ \vec{\nabla}' \times \vec{J} \right]_{\text{ret}}. \quad (20)$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials

Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction


Non-relativistic applications

## 2.2- Retarded electromagnetic field

- The expressions for the retarded fields Eqs. 19 and 21 can also be written in a form that connects directly with the static expressions (tarea):

$$\vec{E}(\vec{x}, t) = \frac{1}{4\pi\epsilon_0} \int d^3x' \left\{ \frac{\hat{R}}{R^2} [\rho(\vec{x}', t')]_{\text{ret}} + \frac{\hat{R}}{cR} \left[ \frac{\partial \rho(\vec{x}', t')}{\partial t} \right]_{\text{ret}} - \frac{1}{c^2 R} \left[ \frac{\partial \vec{J}}{\partial t} \right]_{\text{ret}} \right\} \quad (21)$$

$$\vec{B}(\vec{x}, t) = \frac{\mu_0}{4\pi} \int d^3x' \left\{ \left[ \vec{J}(\vec{x}', t') \right]_{\text{ret}} \times \frac{\hat{R}}{R^2} + \left[ \frac{\partial \vec{J}(\vec{x}', t')}{\partial t} \right]_{\text{ret}} \times \frac{\hat{R}}{cR} \right\} \quad (22)$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials

Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications



## 3.1- Wave zone Wave zone

- We now consider harmonic sources (the general case can be obtained by superposition of such sources):

$$\begin{aligned}\rho(\vec{x}, t) &= \rho(\vec{x})e^{-i\omega t}, \\ \vec{J}(\vec{x}, t) &= \vec{J}(\vec{x})e^{-i\omega t}.\end{aligned}\tag{23}$$


- We saw that in the presence of sources, the field  $\vec{A}(\vec{x}, t)$  generated in vacuum, and without spatial boundaries, is

$$\vec{A}(\vec{x}, t) = \frac{\mu_0}{4\pi} \int d^3x' \int dt' \frac{\vec{J}(\vec{x}', t')}{|\vec{x} - \vec{x}'|} \delta\left(t' + \frac{|\vec{x} - \vec{x}'|}{c} - t\right).$$

- For harmonic sources,

$$\vec{A}(\vec{x}, t) = e^{-i\omega t} \frac{\mu_0}{4\pi} \int d^3x' \frac{\vec{J}(\vec{x}') e^{ik|\vec{x} - \vec{x}'|}}{|\vec{x} - \vec{x}'|},\tag{24}$$

with  $k = \omega/c$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

## 3.1- Wave zone Wave zone


- We then obtain  $\vec{H}$  and  $\vec{E}$  with

$$\vec{H} = \frac{1}{\mu_0} \vec{\nabla} \times \vec{A}, \quad (25)$$

and Faraday's law,

$$\vec{E} = \frac{i}{k} \sqrt{\frac{\mu_0}{\epsilon_0}} \vec{\nabla} \times \vec{H}. \quad (26)$$

- We now consider sources confined inside a region whose maximum extension is  $d$ , and that contains the origin. If  $d \ll \lambda$ , there are 3 regions of interest:
  - The near zone, with  $d < r \ll \lambda$ , where  $e^{ik|\vec{x}-\vec{x}'|} \sim 1$  and we recover the static potentials except for harmonic oscillation,  $\vec{A}(\vec{x}, t) = \vec{A}(\vec{x})e^{-i\omega t}$ .
  - The intermediate zone with  $d \ll r \sim \lambda$ .
  - The far zone, with  $d \ll r$ .



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction


Non-relativistic applications

## 3.1- Wave zone Wave zone

- In the far zone, with  $d \ll r$ ,  $|\vec{x} - \vec{x}'| \approx r - \hat{n} \cdot \vec{x}'$ , where  $\hat{n} = \vec{x}/r \Rightarrow$

$$\lim_{kr \rightarrow \infty} \vec{A}(\vec{x}) = \frac{\mu_0}{4\pi} \frac{e^{ikr}}{r} \int \vec{J}(\vec{x}') e^{-ik\hat{n} \cdot \vec{x}'} d^3x'. \quad (27)$$

- We see that  $\vec{A}(x, t) = \vec{A}(\vec{x})e^{-i\omega t}$  represents a spherical wave travelling outwards.
- In addition, (tarea) using Eqs 25 and 26 we also see that  $\vec{E}$  and  $\vec{H}$  also form spherical transverse waves (orthogonal to  $\hat{n}$ ).
- The far zone thus corresponds to the *radiation zone*, also called *wave zone*.



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications


## 3.2- Dipolar radiation Dipolar radiation

- We now use  $d \ll \lambda$  to simplify  $\vec{A}$  in the wave zone. The integrand in Eq. 27 can be expanded in powers of  $-ik\hat{n} \cdot \vec{x}'$ , using

$$e^{-ik\hat{n} \cdot \vec{x}'} = \sum_{n=0}^{\infty} \frac{(-ik)^n}{n!} (\hat{n} \cdot \vec{x}')^n.$$

- Therefore,

$$\vec{A}(\vec{x}) = \frac{\mu_0}{4\pi} \frac{e^{ikr}}{r} \sum_{n=0}^{\infty} \frac{(-ik)^n}{n!} \int \vec{J}(\vec{x}') (\hat{n} \cdot \vec{x}')^n d^3x'. \quad (28)$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications

## 3.2- Dipolar radiation Dipolar radiation

- For  $n = 0$ , which is the dominant term in the expansion in  $k\hat{n} \cdot \vec{x}'$ , we get:


$$\vec{A}(\vec{x}) = \frac{\mu_0}{4\pi} \frac{e^{ikr}}{r} \int \vec{J}(\vec{x}') d^3x'. \quad (29)$$

- Using the continuity equation,  $\frac{\partial \rho}{\partial t} + \vec{\nabla} \cdot \vec{J} = 0$ , we have  $i\omega\rho = \vec{\nabla} \cdot \vec{J}$ .
- Therefore (tarea):

$$\int \vec{J}(\vec{x}') d^3x' = - \int \vec{x}' (\vec{\nabla}' \cdot \vec{J}) d^3x' = -i\omega \int \vec{x}' \rho(\vec{x}') d^3x'. \quad (30)$$

- Finally,

$$\vec{A}(\vec{x}) = \frac{-i\mu_0\omega}{4\pi} \vec{p} \frac{e^{ikr}}{r}, \quad \text{with} \quad \underbrace{\vec{p} = \int \vec{x}' \rho(\vec{x}') d^3x'}_{\text{electric dipole}}. \quad (31)$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials

Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications

## 3.2- Dipolar radiation Dipolar radiation

- We now calculate the  $\vec{E}$  and  $\vec{H}$  fields:

$$\begin{aligned}\vec{H} &= \frac{ck^2}{4\pi} (\hat{n} \times \vec{p}) \frac{e^{ikr}}{r}, \\ \vec{E} &= \sqrt{\mu_o \epsilon_o} \vec{H} \times \hat{n},\end{aligned}\tag{32}$$


where we see that electric dipole radiation is linearly polarized.

- The power emitted in direction  $\hat{n}$  can be written with  $dP = r^2 d\Omega \hat{n} \cdot \vec{S} \Rightarrow$ ,

$$\begin{aligned}\frac{dP}{d\Omega} &= \frac{1}{2} \Re \left[ r^2 \hat{n} \cdot (\vec{E} \times \vec{H}^*) \right], \\ &= \frac{c^2}{32\pi^2} \sqrt{\frac{\mu_o}{\epsilon_o}} k^4 |\vec{p}|^2 \sin^2(\theta).\end{aligned}\tag{33}$$

- The total power is

$$P = \frac{c^2 k^4}{12\pi} |\vec{p}|^2.\tag{34}$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications



### 3.3-Magnetic dipole and electric quadrupole radiation

- The next term in the expansion of  $e^{-ik\hat{n}\cdot\vec{x}'} = \sum_{n=0}^{\infty} \frac{(-ik)^n}{n!} (\hat{n} \cdot \vec{x}')^n$  corresponds to  $n = 1$ .
- Eq. 28 gives:


$$\vec{A}(\vec{x}) = \frac{\mu_0}{4\pi} \frac{e^{ikr}}{r} \left( \frac{1}{r} - ik \right) \int \vec{J}(\vec{x}') (\hat{n} \cdot \vec{x}') d^3x'. \quad (35)$$

- This term originates *magnetic dipole* and *electric quadrupole* contributions. To see this, we separate the integrand:

$$\vec{J}(\vec{x}') (\hat{n} \cdot \vec{x}') = \underbrace{\frac{1}{2} \left[ (\hat{n} \cdot \vec{x}') \vec{J} + (\hat{n} \cdot \vec{J}) \vec{x}' \right]}_A + \underbrace{\frac{1}{2} (\vec{x}' \times \vec{J}) \times \hat{n}}_B. \quad (36)$$

- We first consider the contribution of part *B* and identify the magnetization  $\vec{\mathcal{M}}$ ,

$$\vec{\mathcal{M}} = \frac{1}{2} (\vec{x} \times \vec{J}). \quad (37)$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

### 3.3-Magnetic dipole and electric quadrupole radiation

- Then, for the  $B$  part,

$$\vec{A}(\vec{x}) = \frac{ik\mu_o}{4\pi}(\hat{n} \times \vec{m}) \frac{e^{ikr}}{r} \left(1 - \frac{1}{ikr}\right), \quad \text{with} \quad (38)$$

$$\vec{m} = \int \vec{\mathcal{M}} d^3x. \quad (39)$$


- In the radiation zone,  $kr \gg 1$ , we obtain:

$$\vec{A}(\vec{x}) = \frac{ik\mu_o}{4\pi}(\hat{n} \times \vec{m}) \frac{e^{ikr}}{r}, \quad (40)$$

$$\vec{E}(\vec{x}) = -\frac{k^2}{4\pi} \sqrt{\frac{\mu_o}{\epsilon_o}} (\hat{n} \times \vec{m}) \frac{e^{ikr}}{r}, \quad (41)$$

$$\vec{H}(\vec{x}) = -\sqrt{\frac{\epsilon_o}{\mu_o}} (\vec{E} \times \hat{n}). \quad (42)$$

- This contribution is called magnetic dipole radiation.



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction


Non-relativistic applications

### 3.3-Magnetic dipole and electric quadrupole radiation

- For the  $A$  part in the contribution from  $n = 1$ , after standard handling we get:

$$A = \frac{i\omega}{2} \int \vec{x}' (\hat{n} \cdot \vec{x}') \rho(\vec{x}') d^3x', \quad (43)$$

which represents order 2 moments of  $\rho(\vec{x}')$ , i.e. an electric quadrupole contribution, which we will not develop.



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation

Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula  
Radiation reaction

Non-relativistic applications


## 4.1- Liénard-Wiechert potentials

- Consider a particle with charge  $q$  and trajectory  $\vec{r}(t)$ , with velocity  $\vec{u}(t)$ .
- We can apply the retarded solution with source terms  $\rho(\vec{x}, t) = q\delta(\vec{x} - \vec{r}(t))$ , and  $\vec{j}(\vec{x}, t) = \rho(\vec{x}, t)\vec{u}(t)$ , and thus obtain the resulting potentials (tarea):

$$\Phi(\vec{x}, t) = \frac{1}{4\pi\epsilon_0} \left[ \frac{q}{\left(1 - \hat{n}(t') \cdot \vec{\beta}(t')\right) R(t')} \right]_{\text{ret}}, \quad (44)$$

$$\vec{A}(\vec{x}, t) = \frac{\mu_0}{4\pi} \left[ \frac{q\vec{u}}{\left(1 - \hat{n}(t') \cdot \vec{\beta}(t')\right) R(t')} \right]_{\text{ret}}, \quad (45)$$

where  $\vec{R}(t') = \vec{x} - \vec{r}(t')$ ,  $R = |\vec{R}|$ ,  $\hat{n}(t') = \frac{\vec{R}}{R}$ , and  $\vec{\beta}(t') = \frac{\vec{u}(t')}{c}$ . These are the Liénard-Wiechert (L.-W.) potentials.



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula  
Radiation reaction

Non-relativistic applications


## 4.1- Liénard-Wiechert potentials

- We can also calculate the corresponding electromagnetic field (tarea):

$$\vec{E}(\vec{x}, t) = \underbrace{\left[ \frac{q}{4\pi\epsilon_0} \frac{(1 - \beta^2)(\hat{n} - \vec{\beta})}{R^2(1 - \hat{n} \cdot \vec{\beta})^3} \right]_{\text{ret}}}_{E_{\text{vel}}} + \underbrace{\left[ \frac{q}{4\pi\epsilon_0} \frac{(\hat{n} \times ((\hat{n} - \vec{\beta}) \times \dot{\vec{\beta}}))}{cR(1 - \hat{n} \cdot \vec{\beta})^3} \right]_{\text{ret}}}_{E_{\text{rad}}} \quad (46)$$

with

$$\vec{B}(\vec{x}, t) = \frac{1}{c} \hat{n} \times \vec{E}(\vec{x}, t). \quad (47)$$



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula  
Radiation reaction


Non-relativistic applications

## 4.1- Liénard-Wiechert potentials

- We see that far away from the particle, the term labelled  $E_{\text{rad}}$  will eventually dominate. In fact, for a Fourier component, or for harmonic motion with  $\vec{r}(t) \propto \exp(i\omega t)$ , we find that (tarea)

$$\frac{E_{\text{rad}}}{E_{\text{vel}}} = \beta \frac{R}{\lambda}, \quad (48)$$

and we see that the radiation term dominates if  $R \gg \lambda/\beta$ , sometimes also called the *far zone*.



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula  
Radiation reaction

Non-relativistic applications



# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications


## 4.2- Larmor Formula

- In the far zone, with  $R \rightarrow \infty$ , and in the Galilean limit ( $\beta \ll 1$ ), the power radiated per unit solid angle is (tarea):

$$\frac{dP}{d\Omega} = \frac{\mu_0}{16\pi^2} q^2 a^2 \sin^2(\theta), \quad (49)$$

in which  $a = |\dot{\vec{u}}|$  and  $\theta$  is the angle between  $\hat{n}$  and  $\vec{a}$ . This is the Larmor formula.

- By applying Eq. 49 to the case of an harmonically oscillating charge with dipole  $\vec{p} = q\vec{r}_0 \exp(i\omega t)$ , we can recover Eq. 33 (tarea).
- We therefore conclude that the *wave zone* as defined in Sec. 1 matches the Galilean limit (see the discussion on the dipole approximation in Sec. 3.3 of Rybicki & Lightman).



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

## 4.3- Radiation reaction

- Let's consider a periodic system, such that its mechanical state is identical between times  $t_1$  and  $t_2$ .
- Still in the Galilean limit, the total energy radiated by the charge between  $t_1$  and  $t_2$  is

$$W = \frac{\mu_0 q^2}{6\pi c} \int_{t_1}^{t_2} a^2 dt. \quad (50)$$


- Energy conservation requires the existence of a 'radiation reaction' force  $\vec{F}_{\text{rad}}$ , such that  $W$  is extracted from the particle's kinetic energy,

$$\int_{t_1}^{t_2} dt \vec{F}_{\text{rad}} \cdot \vec{u} = -W. \quad (51)$$

- One expression for the radiation reaction is the Abraham-Lorentz formula,

$$\vec{F}_{\text{rad}} = \frac{\mu_0 q^2}{6\pi c} \dot{\vec{a}}. \quad (52)$$

We can confirm that Eq. 52 indeeds fullfills Eq. 51 (tarea).



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula

Radiation reaction

Non-relativistic applications

# Outline

## 1 Green function for the wave equation

## 2 Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field


## 3 Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation

## 4 Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

## 5 Non-relativistic applications



Green function for the wave equation

Retarded Potentials

Application of the Green function to the electrodynamic potentials  
Retarded electromagnetic field

Multipolar Radiation

Wave zone  
Dipolar radiation  
Magnetic dipole and electric quadrupole radiation


Radiation from a single charge

Liénard-Wiechert potentials  
Larmor Formula  
Radiation reaction

Non-relativistic applications

## 5- Non-relativistic applications

- Thomson scattering (Rybicki & Lightman Sec. 3.4)
- Harmonically bound particles (Rybicki & Lightman Sec. 3.6)



Green function for the wave equation

### Retarded Potentials

Application of the Green function to the electrodynamic potentials

Retarded electromagnetic field

### Multipolar Radiation

Wave zone

Dipolar radiation

Magnetic dipole and electric quadrupole radiation

### Radiation from a single charge

Liénard-Wiechert potentials

Larmor Formula

Radiation reaction

### Non-relativistic applications