



ISM

Simon Casassus Astronomía, Universidad de Chile


<http://www.das.uchile.cl/~simon>

- I Introduction: Observations of the ISM
- II Microscopic Processes.
- III Astrophysics of Gaseous Nebulae.
- IV Interstellar Dust.
- V Dynamics of the ISM.
- VI Selectec topics: protoplanetary disks, planetary nebulae, SNRs.

Part IV

Interstellar dust

- 1 Extinction
- 2 IR emission from dust
- 3 VSGs
- 4 The cycle of interstellar dust




Extinction

IR emission from dust

VSGs

The cycle of interstellar dust



References:

Draine, B.T., 2003, "Astrophysics of Dust in Cold Clouds", astro-ph/0304488
Draine, B.T., 2004, ARA&A, 41, 241

- $A_\lambda \equiv -2.5 \log_{10} I_{\text{obs}}/I_{\text{int}}$.
- $I_{\text{obs}} = \exp(-\tau)I_{\text{int}} \implies A_\lambda = 2.5\tau/\ln(10) \approx 1.086\tau(\lambda)$.
- A_λ is measured by comparing magnitudes for star with the same spectral type, or else using recombination lines.
- $R_V = A_V/E(B - V) = A_V/(B - V|_{\text{obs}} - B - V|_{\text{int}}) = A_V/(A_B - A_V)$

Extinction

IR emission from dust

VSGs

The cycle of interstellar dust


Extinction

- grains $\ll \lambda$: Rayleigh-scattering gives $A_\lambda \propto \lambda^{-4}$, $R_V \approx 1.2$.
- grains $\gg \lambda$: gray extinction, independent of λ , $R_V \rightarrow \infty$.
- Parametrisation in terms of single parameter R_V OK for $\lambda > 3030 \text{ \AA}$:

$$A(\lambda), R_V = a(\lambda) + b(\lambda)/R_V,$$

where $a(\lambda)$ and $b(\lambda)$ are power laws, set by fitting the data (Cardelli et al., 1989, ApJ, 345, 245).

- For $\lambda < 3030 \text{ \AA}$, use $A_\lambda \approx f(\lambda; R_V, \{a_i\})$ where $\{a_i\}$ characterise the UV ‘bump’.
- In the IR, between 0.9 and $6 \mu\text{m}$, A_λ is \pm universal: $A_\lambda = 2.4 \lambda^{-1.75}$ (Draine 1989, quoting ‘Infrared Astronomy’, Tokunaga in *Astrophysical Quantities*).




Extinction

IR emission from dust

VSGs


The cycle of interstellar dust

Extinction



B. T. Draine 2003.04.03

ISM




Extinction

IR emission from dust

VSGs


The cycle of interstellar dust

Extinction



B. T. Draine 2003.04.25

ISM




Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

- Galactic bulge: $R_V = 1.8 - 2.5$ (Udalski 2002)
- LMC: $R_V = 3.1$ (Udalski 2002)
- Cirrus clouds (far from the galactic plane): $R_V = 2$ (Guhathakurta 1999).
- Typical value for the local ISM : $R_V = 3.1 \pm 2!$? (Cardelli et al 1989)



Extinction


IR emission from dust

VSGs

The cycle of interstellar dust


$$N_H \leftrightarrow A_V^1$$

Using H I Ly α and H₂ absorption lines to measure the column of protons to nearby stars gives $N_H/E(B - V) = 5.8 \cdot 10^{21} \text{ cm}^{-2} \text{ mag}^{-1}$, or $N_H/A_V = 1.87 \cdot 10^{21} \text{ cm}^{-2} \text{ mag}^{-1}$, for $R_V = 3.1$. But $N_H \leftrightarrow A_V$ varies with sightlines:



$A_V(N_H)$ follows Z in the Magellanic Clouds: 40% of Milky Way extinction in LMC, 10% in SMC.

¹ojo: A_V vs $1/R_V \propto 1/A_V$




Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

A_λ over 6-8 μm



ISM



Extinction



IR emission from dust

VSGs

The cycle of interstellar dust

A_λ over 6-8 μm

Lutz et al. (1996, A&A, 315, L269) find excess extinction assuming Case B for H recombination lines in the *ISO*+*SWS* spectrum.




Extinction

IR emission from dust

VSGs


The cycle of interstellar dust

A_λ over 6-8 μm



B. T. Draine 2002.12.05

ISM




Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

Uncertainties in A_λ



Dwek (2004, ApJL, 611, 109) proposes that the excess extinction at $\sim 6 \mu\text{m}$ is due to metallic needles (ref. Hoyle & Wickramasinghe).

short dash:

silicates, **long dash:**

metallic needles, **dots:**


graphite, **dot-dash:**

PAHs, **thin solid:**

standard extinction.

$$\tau(\lambda) = \sum_j \int_0^L ds \left[\int_{a_{\min,j}}^{a_{\max,j}} da f_j(a) \kappa_j(\lambda, a) \rho_{d,j}(a, s) \right], \quad \text{with}$$

$$\int_{a_{\min,j}}^{a_{\max,j}} f_j(a) da = 1.$$



Extinction



IR emission from dust

VSGs

The cycle of interstellar dust

The UV bump: VSGs

- The UV bump matches a Lorentzian (in truth a 'Drude' profile).
- The fitting of a Lorentzian allows inferring $n_X f_X / n_H = 9.3 \cdot 10^{-6}$, and since the oscillator strength of the absorbing specie cannot be in excess of ~ 1 , $f_X \lesssim 0.5$, one requires $n_X / n_H \gtrsim 4 \cdot 10^{-5}$, allowing to consider compounds of $\{C, N, O, Mg, Si, Fe\}$.
- The centre of the bump is constant at 2175\AA , although the width of the bump varies with $1 \sigma \approx 6\%$.





Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

- A ground state transition in the delocalized electrons of the hexagonal benzene rings lies precisely at 2175\AA .
- Graphite, with $f = 0.16$ could explain λ_o , and the absorption strength with $C/H = 5.8 \cdot 10^{-5}$ (OK for cosmic abundances), but not the FWHM variations.
- Variations in the mixture of different PAHs molecules are required to explain the *FWHM* variations.



Extinction


IR emission from dust

VSGs

The cycle of interstellar dust

Silicate extinction

- The $10\mu\text{m}$ silicate feature is seen in absorption when the screen is colder than the background illuminating source. It is seen in emission in M stars ($C/O > 1$), or in planetary nebulae with $C/O > 1$.
- Stretching mode of Si-O at $9.7\ \mu\text{m}$.
- Properties of silicates vary substantially with environment.
- The extinction curve at $10\mu\text{m}$ is smooth and shows no fine structure, in marked contrast with the crystalline silicate spectra \Rightarrow interstellar silicates are amorphous.
- But there is a significant fraction of crystallinity in circumstellart disks and AGB (M) star outflows.
- Bowey et al. propose that up to 60% of silicate mass in dense dark cloud is crystalline (criticised).







Extinction

IR emission from dust

VSGs

The cycle of interstellar dust





Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

Silicate extinction



Extinction



IR emission from dust

VSGs

The cycle of interstellar dust

Variations in the silicate profiles³

ISM



Extinction



IR emission from dust

VSGs


The cycle of interstellar dust

³Van Dischoeck, 2004, AR&AA

Absence of crystalline silicates in the diffuse ISM⁴



⇒ Crystalline silicates represent $\sim 0.2 \pm 0.2$ by mass. Since the silicates in the AGB star outflows have high fractions of crystallinity, there must be substantial interstellar processing of dust in the ISM.



Extinction




IR emission from dust

VSGs

The cycle of interstellar dust

⁴Kemper et al. ApJ 2004, 609, 826

Ice absorption



Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

IR emission from dust

If the grains are in steady state, their internal energy and temperature are constant, and the emergent spectrum from a homogeneous cloud is

$$I_\nu = B_\nu(T_d) [1 - \exp(-\tau_d)],$$

with $\tau = \kappa(\lambda) \int \rho ds$, and


$$\kappa(\lambda) = \frac{1}{1.4n_H m_H} \int da \frac{dn}{da} C_{\text{abs}}(a, \lambda),$$

where ρ is the mass density (with gas), and $n(a)$ is the number density of grains with radii $< a$. Thus the column of mass is

$$\int \rho ds \approx \frac{1}{\kappa(\lambda)} \frac{I_\nu}{B_\nu(T_d)}.$$

The opacities κ and cross sections C_{abs} are tabulated in the web page of B. Draine:

<http://www.astro.princeton.edu/~draine/dust/dust.html>.



Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

IR emission from dust

The power absorbed by a grain is


$$\left. \frac{dE}{dt} \right|_{\text{abs}} = \int_0^{\infty} d\nu u_{\nu} c C_{\text{abs}}(\nu),$$

and emitted power is

$$\left. \frac{dE}{dt} \right|_{\text{rad}} = \int_0^{\infty} d\nu 4\pi B_{\nu} C_{\text{abs}}(\nu).$$

We can express T_s , the steady state temperature, from $\left. \frac{dE}{dt} \right|_{\text{abs}} = \left. \frac{dE}{dt} \right|_{\text{rad}}$, **under the assumption that collisions do not contribute to the thermal balance.**

But in reality E , the internal energy of the grain, and T , its temperature, vary stochastically, and the fluctuations are more pronounced when $h\nu \sim E$, as is the case for very small grains.





Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

Feature at 3.4 μm : C-H aliphatic stretch (carbon chains)



Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

Other dust features

- Diamonds are seen in two Herbig AeBe stars and one AGB star, through emission bands at 3.4 and 3.5 μm due to hydrogenated surface impurities on an otherwise regular carbon-crystal lattice.
- SiC (silicon carbide) is seen towards C stars and PNe.
- The Extended Red Emission, or a broad featureless continuum at 6000 to 8000 \AA , is seen towards all lines of sight (i.e. cirrus and dark clouds, H II regions, PNe and galaxies in general). The carriers of ERE have yet to be identified.
- Fullerenes (Kroto et al. 1985, Nat, 318, 162) in post-AGB stars and PNe, e.g. Bernard-Salas et al., 2012, ApJ, 757, 41B

ISM




Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

- The VSGs are necessary to explain the variations in the UV bump of the extinction curve.
- Andriesse (1978) postulated the VSGs to explain the variation of the dust temperature in a cloud near M17. T_{colour} in the NIR and FIR are 150K y 38K, respectively, while the NIR and FIR morphologies are practically identical \rightarrow the two temperatures are mixed, and it is necessary to postulate the temperature excursions of VSGs (which were called Platt particles at the time).
- Sellgren et al. (1983) found that the NIR emission in reflection nebula is extended and consists at least of one emission band at $3.3 \mu\text{m}$ as well as a continuum with $T_{\text{colour}} = 1000 \text{ K}$, with minimal variations with distance from the central star. Sellgren (1984, ApJ, 277, 623) noted that if the dust was in radiative equilibrium with the star, its temperature should decrease rapidly with distance, and so postulated the VSGs.




Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

- The $12\ \mu\text{m}$ and $25\ \mu\text{m}$ emission from Galactic cirrus, discovered by *IRAS*, is much higher than that expected from large grains in equilibrium with the local radiation field (with $T \sim 18 - 25\ \text{K}$, Boulanger & Perault 1988).
- Spinning dust?




Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

Stochastic heating of VSGs



A grain with radius 5 \AA (i.e. a PAH) receives 1 UV photon (10 eV) per year in the radiation field of the solar neighbourhood. When absorption occurs, the dust grain temperature may rise up to 1000 K.



Extinction

IR emission from dust

VSGs



The cycle of interstellar dust

The state of the art in calculations of the VSG emissivity is that by Draine & Li (2001, ApJ, 551, 807), who estimated the quantized energy levels of the VSGs and their corresponding transition rates. See also Draine & Anderson (1985, 292, 494), Désert et al. (1986, A&A, 160, 295), Guhathakurtha & Draine (1989, ApJ, 345, 230).

The problem involves modelling grains as discrete systems, where the rate of upwards transitions are

$$T_{l \rightarrow u} = C_{\text{abs}}(E_u - E_l) c \frac{u_E}{E_u - E_l} \Delta E_u,$$

and $T_{u \rightarrow l}$ is obtained by detailed balance.



Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

The probability density $P(E)$ in a discretized grain is then obtained by solving:

$$\frac{d}{dt}P_i = \sum_{j \neq i} T_{ij}P_j - \sum_{j \neq i} T_{ji}P_i.$$


Given $P(E)$, the power density emitted by a single grain is

$$S_\lambda = 4\pi \int dE \frac{dP}{dE} C_{\text{abs}}(\lambda) B_\lambda(T(E)).$$

The temperature of the grain can be defined assuming the energy of the grain is equal to its expectation value, for a vibrational system in contact with a heat bath:

$$\langle E \rangle = \sum_j \frac{\hbar\omega_j}{\exp(\hbar\omega_j/kT) - 1},$$

where the sum extends over all normal modes.



Extinction


IR emission from dust

VSGs

The cycle of interstellar dust

It is reasonable to apply thermodynamics to a single molecule, say coronene ($C_{24}H_{12}$), because it has 10^{22} vibrational states with energies inferior to 1 eV $\Rightarrow 1/T = \partial S/\partial E$, in the canonical ensemble.

The function $P(E)$ depends on each astrophysical situation, but the only calculations available apply to the diffuse ISM.





Extinction

IR emission from dust

VSGs

The cycle of interstellar dust





Extinction

IR emission from dust

VSGs

The cycle of interstellar dust



Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

Evolution of dust: sources

- dust features are observed in red giants.
- the “super-wind” on the AGB phase is very enriched in dust (AGB stars are conspicuous *IRAS* sources).
- M stars (with $O/C > 1$ and classified by the presence of molecular lines such as TiO), show the silicate dust feature at $10\mu\text{m}$.
- C stars (with $C/O > 1$ and classified by the presence of molecular lines such as CH, C_2 y CN), show SiC features and graphite continuum.

The details of the condensation of dust are complex. It is believed that condensation occurs in “detashed shells” of the AGB phase (shielding from the stellar radiation is required to cool the shell and condense elements).




Extinction

IR emission from dust

VSGs


The cycle of interstellar dust

Sources of interstellar dust



CO(1-0) and CO(2-1) in UCamelpardalis (Lindqvist et al. 1998, 351, L1) \rightarrow He-shell flashes modulate the rate of mass loss from $\sim 10^{-7}M_{\odot} \text{ yr}^{-1}$ to $\sim 10^{-5}M_{\odot} \text{ yr}^{-1}$.

ISM




Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

AGB stars



Monnier et al. (2004, ApJ, 605, 436).




Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

Evolution of dust: IRC+10216



Time evolution of the dusty envelope (Weigelt et al. 2002, A&A,392, 131).






Extinction

IR emission from dust

VSGs

The cycle of interstellar dust

Evolution of dust: the PN phase




Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

Evolution of dust: the PN phase and the C/O dichotomy



ISM



Extinction


IR emission from dust

VSGs


The cycle of interstellar dust

Evolution of dust: the PN phase and the C/O dichotomy


18um silicates




11.3um PAH



11.3/18



ISM



Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

Evolution of dust: SNe as dust factories

- Red giants and supergiants are easily identified as sources of interstellar dust. Evolved stars inject dust in the ISM. But does circumstellar dust survive the SN event?
- The sub-mm detection of red-shifted dust from primordial galaxies (Ivison et al. 2000, MNRAS, 315, 209) implies the efficient production of dust on very short timescales, such as can only be provided by type II SNe.
- Isotopic anomalies in meteoritic inclusions favor the production of dust following the SN event (i.e. the SuperNOvaeCONDensates).
- The thermal echo due to radiative heating of the dusty winds from the precursor star (as in SN 1987A, Roche 1989, Nature, 337, 533) allows a measure of the circumstellar dust mass, and thereby a constraint on the precursor spectral type. Such a diagnostic is a lot more difficult from the near-IR spectra (uncertain contribution from VSGs and other excitation mechanisms).

ISM




Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

SN 1987A



The ejecta detected by Bouchet et al. corresponds to the exponentially decaying mid-IR lightcurve. The equatorial ring is currently heated by the collision of the supernova blast on circumstellar shell.




Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

SN 2002hh: first IR echo since SN 1987A.



- (a, b) SINGS IRAC 5.8 and 8.0 μm images of a 30"x29" region around SN 2002hh (1.1" per pixel), obtained on 2004 June 10.
- (c) SINGS MIPS 24 μm image of the same region (2.6" per pixel), obtained on 2004 July 9.
- (d) Gemini North Michelle 11.2 μm image of SN 2002hh (0.099"), obtained on 2004 September 26.




Extinction

IR emission from dust


VSGs

The cycle of interstellar dust

SN 2002hh: first IR echo since SN 1987A.



The modelling of the radiatively heated dust in SN 2002hh yields a total dust mass of $0.10\text{--}0.15 M_{\odot}$, giving a progenitor mass of $\sim 10\text{--}15 M_{\odot}$. The parameters of the dust shell are reminiscent of those around Galactic LBVs: a radius of $0.5\text{--}1$ pc, dust masses of $0.1\text{--}0.3 M_{\odot}$.



Extinction


IR emission from dust

VSGs

The cycle of interstellar dust

SN 2002hh.

- The IR luminosity of SN 2002hh at day 600 is $\sim 10^7 L_{\odot}$, much higher than that expected for the SN ejecta.
- Hence the observed mid-IR emission is due to flash-heated dust, a kind of “thermal echo”.
- The thermal echo interpretation, rather than the condensation of dust in the SN ejecta, is also supported by the absence of line asymmetries in SN 2002hh for the same epoch (Clayton & Welsh 2004, AAS meeting).
- By comparison, SN 1987A scaled to the distance of SN 2002hh would have a factor of 35 less mid-IR flux at the same epoch.



Extinction


IR emission from dust

VSGs

The cycle of interstellar dust

Dust condensation and emission line asymmetry.

- Among the discoveries in SN 1987A, observed around day 600 but largely unexploited, is the asymmetry of optical emission lines generated by dust internal to the ejecta (Danziger et al. 1989, IAUC 4746). Dust condenses during the cooling related to the violent expansion, after day ~ 300 . Dust absorption extinguishes the red-shifted side of the metal-rich ejecta, thereby blue-shifting the observed profiles. Scattering off dust in an expanding nebula should also generate a red tail in the emission line profiles.




Extinction

IR emission from dust

VSGs


The cycle of interstellar dust

Dust condensation in Cas A.



Rho et al. 2008.

ISM




Extinction

IR emission from dust


VSGs

The cycle of interstellar dust


Evolution of dust: Condensation in hostile environments



Interacting Binary Wind Model
of Spiral Outflow Around WR 104



ISM



Extinction


IR emission from dust

VSGs

The cycle of interstellar dust

Evolution of dust: destruction of dust and time scales⁵

- Presolar grains in meteorites (e.g. in Murchison) show isotopic anomalies associated to stardust. But this does not imply that the bulk of the dust in the ISM is indeed stardust.
- ISM dust is destroyed in SN shock waves (Tielens et al. 1994, ApJ, 431, 321), by sputtering from heavy ions in hot gas, by grain-grain collisions leading to grain vaporization and fragmentation, and by UV exposure, which depletes the VSGs.
- *Models* of the evolution of dust grains for the various phases of the ISM *show* that the time scale for the residence of one atom in a given grain is $\tau_{\text{dest}} \sim 3 \cdot 10^8$ yr.
- The rate of star formation in our Galaxy is $dM/dt \sim 5 M_{\odot} \text{ yr}^{-1}$, and the total mass of the ISM is $M_{\text{ism}} \sim 5 \cdot 10^9 M_{\odot}$. The time scale for the residence of refractory elements in the ISM is thus $\tau_{\text{ism}} \approx M_{\text{ism}}/dMdt \approx 10^9$.



Extinction

IR emission from dust

VSGs


The cycle of interstellar dust

⁵Draine, B.T., 2004, ARA&A, 41, 241

Evolution of dust: destruction of dust and time scales

- Therefore the fraction of τ_{ism} that a given element spends inside a grain is $\tau_{dest}/\tau_{dest} + \tau_{ism} = 0.2$ (why not τ_{dest}/τ_{ism} ?).
→ if no dust is produced in the ISM, then 20% of the refractory elements should be locked in dust.
- But the prediction that 20% of the refractory elements are in dust does not match observations, which give up to 90% for Si → dust is produced in the ISM
- The mass of the Milky Way ISM is uncertain to 1 order of magnitude, and so is the rate of star formation, and therefore the fraction of Si in grains.

ISM



Extinction

IR emission from dust

VSGs

The cycle of interstellar dust